

4<sup>th</sup> International Congress of Turkish Neuroendocrinology Society | ISTANBUL



# Effects of Ramelteon on the contraction of myometrium in rats

Eda COBAN<sup>1</sup>, Ozgur BULMUS<sup>2</sup>

1) Firat University, Faculty of Medicine, Department of Physiology, Elazig, Turkey

2) Balıkesir University, Faculty of Health Sciences, Department of Physiotherapy and Rehabilitation, Balıkesir, Turkey

### Poster # PC40



4<sup>th</sup> International Congress of Turkish Neuroendocrinology Society ISTANBUL



26 28

2020

#### AIM:

- Ramelteon is a melatonin MT1/MT2 receptor agonist, which is used especially in the treatment of sleep disturbanc
- In this study, we aimed to investigate the effects of ramelteon on uterine contractions.

## METHODS:

- Sprague-Dawley female rats were used in the study (n=7).
- Longitudinal myometrium strips (12mmx2mmx1mm) were placed in an isolated organ bath by applying 1 g of tension in krebs solution, and isometric contractions were recorded.
- Following the regulation of spontaneous contractions, 0.5 μM and 1 μM doses of the ramelteon were administered as a double dose at 60 min intervals and monitored for 30 minutes.
- Statistical analyses were performed using the Paired Sample t-Test.



## 4<sup>th</sup> International Congress of Turkish Neuroendocrinology Society ISTANBUL



## **RESULTS**:



- Ramelteon known as melatonin agonist decreases the frequency value by 83.1%, peak to peak (p-p) value by 90% and amplitude value by 97% after the application of 0.5 μM dose.
- In ramelteon treatment at 1 μM dose, it decreased the frequency value by 90%, p-p value by 97% and area value by 100% (statistically, p <0.001).</li>



4<sup>th</sup> International Congress of Turkish Neuroendocrinology Society ISTANBUL



# CONCLUSION:

- It is thought that ramelteon, which is used in the treatment of sleep disorder and can be used in pregnancy, shows its inhibitory activity on uterine contractions over MT1/MT2 receptors such as melatonin.
- Ramelteon is recommended for use during pregnancy, taking into account its inhibitory effect on uterine contractions.